

Mechanism of Cu-Catalyzed Aryl Boronic Acid Halodeboronation Using Electrophilic Halogen: Development of a Base-Catalyzed Iododeboronation for Radiolabeling Applications

John J. Molloy,[†] Kerry M. O'Rourke,[‡] Carolina P. Frias,[§] Nikki L. Sloan,[‡] Matthew J. West,[†] Sally L. Pimlott,^{||} Andrew Sutherland,^{*,‡,||} and Allan J. B. Watson^{*,†,||}


[†]EaStCHEM, School of Chemistry, University of St Andrews, North Haugh, St Andrews, Fife, KY16 9ST, U.K.

[‡]WestCHEM, School of Chemistry, University of Glasgow, Joseph Black Building, University Avenue, Glasgow, G12 8QQ, U.K.

[§]WestCHEM, Department of Pure and Applied Chemistry, University of Strathclyde, 295 Cathedral Street, Glasgow, G1 1XL, U.K.

^{||}West of Scotland PET Centre, Greater Glasgow and Clyde NHS Trust, Glasgow, G12 0YN, U.K.

S Supporting Information



ABSTRACT: An investigation into the mechanism of Cu-catalyzed aryl boronic acid halodeboronation using electrophilic halogen reagents is reported. Evidence is provided to show that this takes place via a boronate-driven *ipso*-substitution pathway and that Cu is not required for these processes to operate: general Lewis base catalysis is operational. This in turn allows the rational development of a general, simple, and effective base-catalyzed halodeboronation that is amenable to the preparation of ¹²⁵I-labeled products for SPECT applications.

Halodeboronation of aryl boronic acids is a useful method for the synthesis of aryl halides and has found important applications in the generation of radiolabeled products for *in vivo* imaging and radiotherapy (Scheme 1a).^{1,2} The use of molecular halogen (X₂) and electrophilic halogen (X⁺) sources is very common with several approaches developed, including metal-free (Scheme 1b)³ and metal-promoted methods: Cu-catalysis is particularly prominent in this latter area (Scheme 1c).⁴ Iodo-, bromo-, and chlorodeboronation operate effectively with X⁺ reagents (e.g., halosuccinimide reagents), and while fluorodeboronation using F⁺ sources has been achieved,^{3b,c,f} the majority of approaches for C–F installation employ alternative catalytic manifolds that utilize F[–].^{2,5}

In terms of metal-free processes, in 1930, Challenger reported the halodeboronation of PhB(OH)₂ using aqueous X₂ as well as with CuX₂, with the latter also proceeding via a pathway involving X₂ produced *in situ* by the known redox of CuX₂ → CuX + X₂.^{3a} In the 1950s, Kuivila established kinetic parameters for specific aryl boronic acid halodeboronation using X₂ in both acidic and basic buffer, proposing an *ipso*-substitution proceeding via a boronate generated *in situ* as the most likely mechanism (Scheme 1d).^{6,7} Boronate-based *ipso*-substitution has also been proposed in halodeboronation reactions of aryl potassium trifluoroborates (ArBF₃K).^{3f,g,i,j}


In contrast, despite a number of variants, the mechanism of Cu-catalyzed halodeboronation reactions using X⁺ reagents remains underdeveloped, mainly relying on plausible constructs with limited empirical support.⁴ Mechanistic proposals have suggested a process via oxidative addition of Cu(I) to a halosuccinimide or derivative, transmetalation of an organoboron to the resulting Cu(III) intermediate, and subsequent reductive elimination (generalized in Scheme 1d).^{4a,b,e} However, there is little evidence to support this proposed mechanism. Accordingly, based on the broad utility of halodeboronation methods, a functional understanding would be valuable to assist in the development and expansion of applications.

Here we provide a mechanistic investigation of arylboronic acid halodeboronation using X⁺ reagents, showing that these reactions proceed via a common base-catalyzed *ipso*-substitution mechanism, and that specific “Cu-catalyzed” reactions do not require Cu to proceed. This allows rational design of a simple KOAc-catalyzed halodeboronation, amenable to the transition metal-free preparation of SPECT radiotracers.

Received: March 16, 2019

Published: March 25, 2019

Scheme 1. Halodeboronation of Boronic Acids and Derivatives: (a) General Reaction; (b) Metal-Free Methods; (c) Cu-Catalyzed Approaches; (d) Proposed Mechanisms^a




^aNXS = N-halosuccinimide.

On the basis of our combined interests in Cu-catalyzed transformations of organoborons⁸ and radiohalogenation of functionalized arenes,⁹ we sought to understand Cu-catalyzed halodeboronation to allow tailoring of specific applications. Accordingly, we initiated our study by evaluating Cu-catalyzed halodeboronation of boronic acids using N-halosuccinimides. A general benchmark reaction was constructed on the basis of previously published methods:⁴ boronic acid **1a** was identified as a workhorse substrate (previously investigated by Kuivila^{7c}) and N-iodosuccinimide (NIS) was selected as a benchmark X⁺ reagent (Scheme 2a). Consistent with literature precedent, using Cu(OAc)_n (n = 1, 2) as a catalyst, the desired iodoarene **2a** was delivered in good yield, with ca. 5% protodeboronation. However, the iododeboronation was evident in the absence of any catalyst, albeit in low yield (30%), consistent with observations by Petasis and Olah (*vide infra*).^{3d} Accordingly, we sought to determine the origin of rate enhancement using Cu.

¹¹B NMR was informative (Scheme 2b). Boronate signals attributed to **4a** were clearly observed in the presence of CuOAc, along with some protodeboronation (production of B(OH)₄[−] (**5**)). **4a** was not observed in the presence of Cu(OAc)₂, with significant protodeboronation to deliver **5** as the main outcome. Protodeboronation in the presence of Cu(II) is consistent with initial observations by Challenger^{3a} and in agreement with additional studies.^{10,11} Control experiments using KOAc and KOH resulted in the formation of the same signal for **4a**, as expected.¹² On the basis of the observed formation of boronate in the presence of CuOAc, the implication of boronate-driven protodeboronation in the presence of Cu(OAc)₂, and that metal-free *ipso*-halodeboronation is well-known,³ we queried whether Cu was required in these processes.

Exposure of **1a** to KOAc delivered **2a** in excellent yield and with a profile commensurate to Cu(OAc)₂ (Scheme 2c). A brief analysis of other bases revealed the process was generally effective in the presence of all Lewis bases capable of forming a boronate. Collectively, these data supported the hypothesis that “Cu-catalyzed” *ipso*-halodeboronation was actually a

Scheme 2. Evidence To Support Halodeboronation via *ipso*-Substitution Pathway



^aReaction at 50 °C.

general base-catalyzed phenomenon and that Cu had no direct role in the catalysis.

An *ipso*-halodeboronation pathway was further supported by straightforward control reactions using specific organoboron reagents with established reactivity profiles (Scheme 2d). If proceeding via boronate, BPin species **1b** would be expected to be less reactive than the equivalent boronic acid **1a** based on the lower propensity to form boronates.¹¹ Exposure of **1b** to NIS gave trace amounts of product **2a** in the presence and absence of catalytic KOAc. As a preformed stoichiometric boronate, the BF₃K species **1c** operated equally effectively in the presence and absence of base as would be expected and consistent with several studies using BF₃K species.^{3f,g,i,j} BMIDA **1d** should not be capable of halodeboronation as this species is not capable of forming a boronate and, while formally zwitterionic, would require the significantly unfavorable loss of a divalent cationic boron leaving group (⁺BR₂).¹³

Indeed, **1d** did not react with NIS either at rt or 50 °C. However, the related BIDA compound **1e**, lacks the N-Me unit of BMIDA and is therefore capable of proton loss from the ligating N–H and liberation of a neutral (BR₃) leaving group (not possible for BMIDA). On evaluation, **1e** did deliver the expected product in up to moderate yield and was again accelerated by KOAc. Collectively, these data were consistent with a boronate *ipso*-substitution pathway.

Additional support for *ipso*-substitution came from a straightforward competition experiment (Scheme 2e). Exposing a mixture of boronic acids **1m** and **1s** to 20 mol % KOAc delivered a mixture of boronates **4m** and **4s**, favoring the more Lewis acidic boronate **4m** (ca. 2:1 **4m**:**4s**). However, since the *ipso*-substitution is rate limiting⁷ and boronate formation/equilibration precedes this event, the 1:1 competition experiment should yield **2s** as the main product. Indeed, a ratio of ca. 9:1 **2s**:**2m** was observed in this experiment.


Regarding reactivity in the absence of base, this has been previously observed by Petasis and Olah.^{3d} This is likely to proceed via the well-known boronic acid autoionization initially reported by Lorand and Edwards,¹⁴ with yields increasing with temperature in agreement with this mechanism.

Kuivila proposed a boronate-driven *ipso*-substitution pathway for halodeboronation using X⁺ reagents.⁷ However, the absence of NMR spectroscopy and access to modern organoboron reagents at the time of study precluded spectroscopic evidence of boronate formation. The body of data presented here strongly supports Kuivila's proposal of boronate-driven *ipso*-substitution.

On the basis of these observations, we proposed that a general and very straightforward base-catalyzed halodeboronation, in line with our SPECT labeling interests, might be realized. Assessing a range of standard variables quickly led to a simple and high yielding cold iododeboronation, which was evaluated on a series of simple substrates (Scheme 3a; see Supporting Information for full details of solvent, temperature, catalyst loading, time, etc.). Using 20 mol % KOAc as catalyst in MeCN at 50 °C allowed smooth halodeboronation in short reaction times, favorable for downstream applications (*vide infra*). Iodo-, bromo-, and chlorodeboronation were all effective; however, in contrast to preceding studies,^{3b,c,f} fluorodeboronation using F⁺ sources was completely ineffective. Consistent with the mechanistic proposal and the above analyses (Scheme 2e), electron-deficient substrates were more sluggish substrates. Some limitations were found with amine substrates; for example, anilines were found to generate several products presumably due to reaction with NIS at the aniline nitrogen.

With the cold method established, we evaluated the compatibility of the method for iododeboronation using ¹²⁵I[−] for imaging applications (Scheme 3b). An optimal transformation was developed using the known method for preparation of I⁺ from I[−], stirring [¹²⁵I]NaI with NCS for 15 min in dimethyl carbonate (DMC).^{9b,15} In accordance with the control studies (Scheme 2a), radio-iododeboronation of **1a** without base gave a radiochemical yield (RCY) of 24% (Scheme 3b). A combination of KOAc (20 mol %) and a reaction temperature of 100 °C, resulted in efficient radioiodination (93%) and in a fast reaction time (1.5 h). This optimized method was applied to a series of boronic acids and found to be tolerant of various functional groups and substitution patterns. In agreement with the cold studies,

Scheme 3. Scope of the Base-Catalyzed Halodeboronation[¶]



[¶]Isolated yields. ^aUsing NBS. ^bUsing NCS. ^cUsing NFSI. ^dNo catalyst and 50 °C. ^e80 °C. ^f10 mol % KOAc. ^gUsing MeCN as cosolvent. ^h50 mol % KOAc.

elevated temperatures were important for complete conversion of electron-deficient substrates.

The utility of the method toward the production of valuable radiolabeled pharmaceuticals was also investigated (Scheme 3b). A di-Boc protected derivative of iobenguane (MIBG) (**9p**), a commercially available radiopharmaceutical used for the SPECT imaging of human norepinephrine transporter-expressing cancers was prepared in 56% RCY, although 50 mol % of KOAc was required for this transformation.¹⁶ The method was also effective for the radioiodination of olaparib

derivative (**9q**), a SPECT imaging agent of the cancer target, poly(ADP-ribose) polymerase-1.¹⁷ Under the standard reaction conditions, this gave **9q** in 67% RCY. It should be noted that the KOAc-catalyzed ¹²⁵I-labeled synthesis of this compound was more efficient than a recently described gold(I)-mediated radio-iododeboronation.^{9d} Finally, iodouracil (**9r**), a potent inhibitor of the anticancer target dihydropyrimidine dehydrogenase¹⁸ and a precursor of uridine-derived SPECT imaging agents,¹⁹ was radioiodinated in 98% RCY after 1 h. In comparison to other approaches, the use of this base-catalyzed method, which avoids transition-metal catalysts, allows a more rapid and operationally simpler approach for the preparation and purification of radioiodinated compounds for imaging applications. These advantages were exemplified with validation of the method for the synthesis and purification of [¹²⁵I]iodouracil (**9r**). Uracil-5-boronic acid was radio-iodinated using [¹²⁵I]NaI (8–10 MBq) at 80 °C for 1 h. After HPLC purification, [¹²⁵I]iodouracil (**9r**) was isolated in 58 ± 1.3% radioactivity yield. The radiochemical purity of **9r** was measured as >99% (*n* = 3), with a molar activity of 0.53 ± 0.047 GBq μmol⁻¹ (*n* = 3).

In summary, we have provided a mechanistic rationale for halodeboronation of aryl boronic acids, demonstrating that the process proceeds via a boronate-driven *ipso*-substitution pathway, providing evidence that specific Cu-catalyzed processes do not require Cu. These observations have allowed the development of a simple base-catalyzed halodeboronation that is amenable to the preparation of ¹²⁵I-labeled products for imaging applications.

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.9b00942.

Experimental procedures, characterization data, copies of spectra (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: andrew.sutherland@glasgow.ac.uk.

*E-mail: aw260@st-andrews.ac.uk.

ORCID

Andrew Sutherland: 0000-0001-7907-5766

Allan J. B. Watson: 0000-0002-1582-4286

Author Contributions

All authors have given approval to the final version of the manuscript.

Funding

EPSRC (EP/R511705/1, EP/K503058/1). University of Glasgow. University of St Andrews. CAPES/Science Without Borders (CAPES/SwB Proc. no. BEX5935/13-0).

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors are grateful to EPSRC (EP/R511705/1 and a studentship to K.M.O., EP/K503058/1) and the University of Glasgow. We thank the University of St Andrews (M.J.W.) and

CAPES/Science Without Borders (C.P.F) for Ph.D. studentships.

■ ABBREVIATIONS

CFS, cesium fluoroxysulfate; DMC, dimethylcarbonate; IDA, iminodiacetoxy; MIDA, *N*-methyliminodiacetoxy; NBS, *N*-bromosuccinimide; NCS, *N*-chlorosuccinimide; NFSI, *N*-fluorobenzenesulfonimide; NIS, *N*-iodosuccinimide; RCY, radiochemical yield; SPECT, single photon emission computed tomography.

■ REFERENCES

- (1) *Boronic Acids: Preparation and Applications in Organic Synthesis, Medicine and Materials*, 2nd ed.; Hall, D., Ed; Wiley VCH: Weinheim, Germany, 2011; Vols. 1 and 2, pp 1–133.
- (2) (a) Preshlock, S.; Tredwell, M.; Gouverneur, V. ¹⁸F-Labeling of Arenes and Heteroarenes for Applications in Positron Emission Tomography. *Chem. Rev.* **2016**, *116*, 719–766. (b) Wilson, T. C.; Cailly, T.; Gouverneur, V. Boron Reagents for Divergent Radiochemistry. *Chem. Soc. Rev.* **2018**, *47*, 6990–7005.
- (3) For examples, see: (a) Ainley, A. D.; Challenger, F. Studies of the Boron-Carbon Linkage. Part I. The Oxidation and Nitration of Phenylboric Acid. *J. Chem. Soc.* **1930**, *0*, 2171–2180. (b) Clough, J. M.; Diorazio, L. J.; Widdowson, D. A. A New Route to Fluoroaromatics from Boronic Acids and Caesium Fluoroxysulfate. *Synlett* **1990**, *1990*, 761–762. (c) Diorazio, L. J.; Widdowson, D. A.; Clough, J. M. A New Synthesis of Aryl Fluorides: The Reaction of Caesium Fluoroxysulfate with Arylboronic Acids and Derivatives. *Tetrahedron* **1992**, *48*, 8073–8088. (d) Thiebes, C.; Prakash, G. K. S.; Petasis, N. A.; Olah, G. A. Mild Preparation of Haloarenes by *Ipso*-Substitution of Arylboronic Acids with *N*-Halosuccinimides. *Synlett* **1998**, *1998*, 141–142. (e) Szumigala, R. H., Jr.; Devine, P. N.; Gauthier, D. R., Jr.; Volante, R. P. Facile Synthesis of 2-Bromo-3-fluorobenzonitrile: An Application and Study of the Halodeboronation of Aryl Boronic Acids. *J. Org. Chem.* **2004**, *69*, 566–569. (f) Cazorla, C.; Métay, E.; Andrioletti, B.; Lemaire, M. Metal-Free Electrophilic Fluorination of Alkyl Trifluoroborates and Boronic Acids. *Tetrahedron Lett.* **2009**, *50*, 3936–3938. (g) Yao, M.-L.; Reddy, M. S.; Yong, L.; Walfish, I.; Blevins, D. W.; Kabalka, G. W. Chemoselective Bromodeboronation of Organotrifluoroborates Using Tetrabutylammonium Tribromide: Application in (*Z*)-Dibromoalkene Syntheses. *Org. Lett.* **2010**, *12*, 700–703. (h) Akula, M. R.; Yao, M.-L.; Kabalka, G. W. Triolborates: Water-Soluble Complexes of Arylboronic Acids as Precursors to Iodoarenes. *Tetrahedron Lett.* **2010**, *51*, 1170–1171. (i) Molander, G. A.; Cavalcanti, L. N. Metal-Free Chlorodeboronation of Organotrifluoroborates. *J. Org. Chem.* **2011**, *76*, 7195–7203. (j) Yao, M.-L.; Kabalka, G. W.; Blevins, D. W.; Reddy, M. S.; Yong, L. Halodeboronation of Organotrifluoroborates Using Tetrabutylammonium Tribromide or Cesium Triiodide. *Tetrahedron* **2012**, *68*, 3738–3743. (k) Niu, L.; Zhang, H.; Yang, H.; Fu, H. Metal-Free Iodination of Arylboronic Acids and the Synthesis of Biaryl Derivatives. *Synlett* **2014**, *25*, 995–1000. (l) Kamei, T.; Ishibashi, A.; Shimada, T. Metal-Free Halogenation of Arylboronate with *N*-Halosuccinimide. *Tetrahedron Lett.* **2014**, *55*, 4245–4247. (m) Tramutola, F.; Chiummiento, L.; Funicello, M.; Lupattelli, P. Practical and Efficient *ipso*-Iodination of Arylboronic Acids via KF/I₂ System. *Tetrahedron Lett.* **2015**, *56*, 1122–1123.
- (4) For examples, see: (a) Savarin, C.; Srogl, J.; Liebeskind, L. S. A Mild Non-Basic Synthesis of Thioethers. The Copper-Catalyzed Coupling of Boronic Acids with *N*-Thio(alkyl, aryl, heteroaryl)imides. *Org. Lett.* **2002**, *4*, 4309–4312. (b) Wu, H.; Hynes, J. H., Jr. Copper-Catalyzed Chlorination of Functionalized Arylboronic Acids. *Org. Lett.* **2010**, *12*, 1192–1195. (c) Zhang, G.; Lv, G.; Li, L.; Chen, F.; Cheng, J. Copper-Catalyzed Halogenation of Arylboronic Acids. *Tetrahedron Lett.* **2011**, *52*, 1993–1995. (d) Ren, Y.-L.; Tian, X.-Z.; Dong, C.; Zhao, S.; Wang, J.; Yan, M.; Qi, X.; Liu, G. A Simple and Effective Copper Catalyst for the Conversion of Arylboronic Acids to Aryl Iodides at Room Temperature. *Catal. Commun.* **2013**, *32*, 15–17.

- (e) He, W.; Zhang, R.; Cai, M. A Highly Efficient Heterogeneous Copper-Catalyzed Chlorodeboronation of Arylboronic Acids Leading to Chlorinated Arenes. *RSC Adv.* **2017**, *7*, 764–770.
- (5) For selected examples, see: (a) Preshlock, S.; Calderwood, S.; Verhoog, S.; Tredwell, M.; Huiban, M.; Hienzsch, A.; Gruber, S.; Wilson, T. C.; Taylor, N. J.; Cailly, T.; Schedler, M.; Collier, T. L.; Passchier, J.; Smits, R.; Mollitor, J.; Hoepping, A.; Mueller, M.; Genicot, C.; Mercier, J.; Gouverneur, V. Enhanced Copper-Mediated ^{18}F -Fluorination of Aryl Boronic Esters Provides Eight Radiotracers for PET Applications. *Chem. Commun.* **2016**, *52*, 8361–8364. (b) Wilson, T. C.; McSweeney, G.; Preshlock, S.; Verhoog, S.; Tredwell, M.; Cailly, T.; Gouverneur, V. Radiosynthesis of SPECT Tracers via a Copper Mediated ^{123}I Iodination of (Hetero)aryl Boron Reagents. *Chem. Commun.* **2016**, *52*, 13277–13280. (c) Taylor, N. J.; Emer, E.; Preshlock, S.; Schedler, M.; Tredwell, M.; Verhoog, S.; Mercier, J.; Genicot, C.; Gouverneur, V. Derisking the Cu-Mediated ^{18}F -Fluorination of Heterocyclic Positron Emission Tomography Radioligands. *J. Am. Chem. Soc.* **2017**, *139*, 8267–8276.
- (6) For a review of metal-free organoboron *ipso*-substitution, see: Zhu, C.; Falck, J. R. Transition Metal-Free *ipso*-Functionalization of Arylboronic Acids and Derivatives. *Adv. Synth. Catal.* **2014**, *356*, 2395–2410.
- (7) (a) Kuivila, H. G.; Easterbrook, E. K. Kinetics of the Brominolysis of Benzenboronic Acid in Aqueous Acetic Acid Solution. *J. Am. Chem. Soc.* **1951**, *73*, 4629–4632. (b) Kuivila, H. G.; Hendrickson, A. R. Electrophilic Displacement Reactions. II. Rates of Brominolysis of Substituted Benzenboronic Acids. *J. Am. Chem. Soc.* **1952**, *74*, 5068–5070. (c) Kuivila, H. G.; Williams, R. M. Electrophilic Displacement Reactions. V. Kinetics of the Iodinolysis of *p*-Methoxybenzenboronic Acid. *J. Am. Chem. Soc.* **1954**, *76*, 2679–2682.
- (8) (a) Vantourout, J. C.; Li, L.; Bendito-Moll, E.; Chabbra, S.; Arrington, K.; Bode, B. E.; Isidro-Llobet, A.; Kowalski, J. A.; Nilson, M.; Wheelhouse, K.; Woodard, J. L.; Xie, S.; Leitch, D. C.; Watson, A. J. B. Mechanistic Insight Enables Practical, Scalable, Room Temperature Chan-Lam *N*-Arylation of *N*-Aryl Sulfonamides. *ACS Catal.* **2018**, *8*, 9560–9566. (b) Vantourout, J. C.; Miras, H. N.; Isidro-Llobet, A.; Sproules, S.; Watson, A. J. B. Spectroscopic Studies of the Chan-Lam Amination: A Mechanism-inspired Solution to Boronic Ester Reactivity. *J. Am. Chem. Soc.* **2017**, *139*, 4769–4779. (c) Vantourout, J. C.; Law, R. P.; Isidro-Llobet, A.; Atkinson, S. J.; Watson, A. J. B. Chan-Evans-Lam Amination of Boronic Acid Pinacol (BPin) Esters: Overcoming the Aryl Amine Problem. *J. Org. Chem.* **2016**, *81*, 3942–3950.
- (9) (a) Cant, A. A.; Champion, S.; Bhalla, R.; Pimlott, S. L.; Sutherland, A. Nickel-Mediated Radioiodination of Aryl and Heteroaryl Bromides: Rapid Synthesis of Tracers for SPECT Imaging. *Angew. Chem., Int. Ed.* **2013**, *52*, 7829–7832. (b) Racys, D. T.; Sharif, S. A. I.; Pimlott, S. L.; Sutherland, A. Silver(I)-Catalyzed Iodination of Arenes: Tuning the Lewis Acidity of *N*-Iodosuccinimide Activation. *J. Org. Chem.* **2016**, *81*, 772–780. (c) Sloan, N. L.; Luthra, S. K.; McRobbie, G.; Pimlott, S. L.; Sutherland, A. A One-Pot Radioiodination of Aryl Amines via Stable Diazonium Salts: Preparation of ^{125}I -Imaging Agents. *Chem. Commun.* **2017**, *53*, 11008–11011. (d) Webster, S.; O'Rourke, K. M.; Fletcher, C.; Pimlott, S. L.; Sutherland, A.; Lee, A.-L. Rapid Iododeboronation with and without Gold Catalysis: Application to Radiolabelling of Arenes. *Chem. - Eur. J.* **2018**, *24*, 937–943.
- (10) Kuivila, H. G.; Reuwer, J. F., Jr.; Mangravite, J. A. Electrophilic Displacement Reactions. XVI. Metal Ion Catalysis in the Protodeboronation of Areneboronic Acids. *J. Am. Chem. Soc.* **1964**, *86*, 2666–2670.
- (11) For selected studies of protodeboronation, see: (a) Cox, P. A.; Reid, M.; Leach, A. G.; Campbell, A. D.; King, E. J.; Lloyd-Jones, G. C. Base-catalyzed Aryl-B(OH) $_2$ Protodeboronation Revisited: From Concerted Proton-Transfer to Liberation of a Transient Aryl Anion. *J. Am. Chem. Soc.* **2017**, *139*, 13156–13165. (b) Cox, P. A.; Leach, A. G.; Campbell, A. D.; Lloyd-Jones, G. C. Protodeboronation of Heteroaromatic, Vinyl and Cyclopropyl Boronic Acids: pH-Rate Profiles, Auto-catalysis and Disproportionation. *J. Am. Chem. Soc.* **2016**, *138*, 9145–9157.
- (12) For example, see: (a) Butters, M.; Harvey, J. N.; Jover, J.; Lennox, A. J. J.; Lloyd-Jones, G. C.; Murray, P. M. Aryl Trifluoroborates in Suzuki–Miyaura Coupling: The Roles of Endogenous Aryl Boronic Acid and Fluoride. *Angew. Chem., Int. Ed.* **2010**, *49*, S156–S160. (b) Molloy, J. J.; Clohessy, T. A.; Irving, C.; Anderson, N. A.; Lloyd-Jones, G. C.; Watson, A. J. B. Chemoselective Brown Oxidation of Aryl Organoboron Systems Enabled by Boronic Acid-selective Phase Transfer. *Chem. Sci.* **2017**, *8*, 1551–155. (c) Molloy, J. J.; Seath, C. P.; West, M. J.; McLaughlin, C.; Fazakerley, N. J.; Kennedy, A. R.; Nelson, D. J.; Watson, A. J. B. Interrogating Pd(II) Anion Metathesis using a Bifunctional Chemical Probe: A Transmetalation Switch. *J. Am. Chem. Soc.* **2018**, *140*, 126–130.
- (13) Matteson, D. S.; Liedtke, J. D. Mechanisms of Deboronobromination. *J. Am. Chem. Soc.* **1965**, *87*, 1526–1531.
- (14) Lorand, J.; Edwards, J. O. Polyol Complexes and Structure of the Benzenboronate Ion. *J. Org. Chem.* **1959**, *24*, 769–774.
- (15) (a) Vankar, Y. D.; Kumaravel, G. *N*-Chlorosuccinimide/Sodium Iodide: A Convenient Source of *N*-Iodosuccinimide. Synthesis of α -Iodo Carbonyl Compounds and *trans*-1,2-Iodoacetates. *Tetrahedron Lett.* **1984**, *25*, 233–236. (b) Youfeng, H.; Coenen, H. H.; Petzold, G.; Stöcklin, G. A Comparative Study of Radioiodination of Simple Aromatic Compounds via *N*-Halosuccinimides and Chloramine-T in TFAA. *J. Labelled Compd. Radiopharm.* **1982**, *19*, 807–819.
- (16) (a) Zhang, H.; Huang, R.; Pillarsetty, N.; Thorek, D. L. J.; Vaidyanathan, G.; Serganova, I.; Blasberg, R. G.; Lewis, J. S. Synthesis and Evaluation of ^{18}F -Labeled Benzylguanidine Analogs for targeting the Human Norepinephrine Transporter. *Eur. J. Nucl. Med. Mol. Imaging* **2014**, *41*, 322–332. (b) Sharp, S. E.; Trout, A. T.; Weiss, B. D.; Gelfand, M. J. MIBG in Neuroblastoma Diagnostic Imaging and Therapy. *Radiographics* **2016**, *36*, 258–278.
- (17) Zmuda, F.; Malviya, G.; Blair, A.; Boyd, M.; Chalmers, A. J.; Sutherland, A.; Pimlott, S. L. Synthesis and Evaluation of a Radioiodinated Tracer with Specificity for Poly(ADP-ribose) Polymerase-1 (PARP-1) in Vivo. *J. Med. Chem.* **2015**, *58*, 8683–8693.
- (18) Dobritzsch, D.; Ricagno, S.; Schneider, G.; Schnackerz, K. D.; Lindqvist, Y. Crystal Structure of the Productive Ternary Complex of Dihydropyrimidine Dehydrogenase with NADPH and 5-Iodouracil. *J. Biol. Chem.* **2002**, *277*, 13155–13166.
- (19) (a) Lan, X.; Liu, Y.; He, Y.; Wu, T.; Zhang, B.; Gao, Z.; An, R.; Zhang, Y. Autoradiography Study and SPECT Imaging of Reporter Gene HSV1-TK Expression in Heart. *Nucl. Med. Biol.* **2010**, *37*, 371–380. (b) Bettgowda, C.; Foss, C. A.; Cheong, I.; Wang, Y.; Diaz, L.; Agrawal, N.; Fox, J.; Dick, J.; Dang, L. H.; Zhou, S.; Kinzler, K. W.; Vogelstein, B.; Pomper, M. G. Imaging Bacterial Infections with Radiolabeled 1-(2'-Deoxy-2'-fluoro- β -D-arabinofuranosyl)-5-iodouracil. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102*, 1145–1150.